

## Tylenol (Acetaminophen) Dosage Chart

**\*\* Tylenol can be given every 4 HOURS \*\* DO NOT give to children under 2 MOS OF AGE \*\***

| Weight (lbs) | Dose (mgs) | Infant Oral Suspension (160mg/5ml) | Children's Oral Suspension (160mg/5ml) | Chewable (80 mg tab) | Jr. Strength (160 mg tab) | Adult tab 325 mg |
|--------------|------------|------------------------------------|--|----------------------|---------------------------|------------------|
| 6-11 lbs     | 40 mg      | 1.25 ml = ¼ tsp                    | ---                                    | ---                  | ---                       | ---              |
| 12-17 lbs    | 80 mg      | 2.5 ml = ½ tsp                     | 2.5 ml = ½ tsp                         | ---                  | ---                       | ---              |
| 18-22 lbs    | 120 mg     | 3.75 ml = ¾ tsp                    | 3.75 ml = ¾ tsp                        | 1 ½ tabs             | ---                       | ---              |
| 23-35 lbs    | 160 mg     | 5 ml = 1 tsp                       | 5 ml = 1 tsp                           | 2 tabs               | 1 tab                     | ---              |
| 36-47 lbs    | 240 mg     |                                    | 7.5 ml = 1½ tsp                        | 3 tabs               | 1½ tab                    | ---              |
| 48-59 lbs    | 320 mg     |                                    | 10 ml = 2 tsp                          | 4 tabs               | 2 tabs                    | 1 tab            |
| 60-71 lbs    | 400 mg     |                                    | 12.5 ml = 2½ tsp                       | 5 tabs               | 2½ tabs                   | 1 tab            |
| 72-95 lbs    | 480 mg     |                                    | 15 ml = 3 tsp                          | 6 tabs               | 3 tabs                    | 1½ tab           |
| 96+ lbs      | 640 mg     |                                    | 20 ml = 4 tsp                          | 8 tabs               | 4 tabs                    | 2 tabs           |

## Motrin & Advil (Ibuprofen) Dosage Chart

**\*\* Ibuprofen can be given every 6 HOURS \*\* DO NOT give to children under 6 MOS OF AGE \*\***

| Weight (lbs) | Dose (mgs) | Infant Oral Suspension (50mg/1.25ml) | Children's Oral Suspension (100mg/5ml) | Chewable (50 mg tab) | Jr. Strength (100 mg tab) | Adult tab 200 mg |
|--------------|------------|--------------------------------------|--|----------------------|---------------------------|------------------|
| 6-11 lbs     | ---        | ---                                  | ---                                    | ---                  | ---                       | ---              |
| 12-17 lbs    | 50 mg      | 1.25 ml=1 drop                       | 2.5 ml = ½ tsp                         | ---                  | ---                       | ---              |
| 18-22 lbs    | 75 mg      | 1.875 ml=1½ drops                    | 3.75 ml = ¾ tsp                        | 1 ½ tabs             | ---                       | ---              |
| 23-35 lbs    | 100 mg     | 2.5 ml=2 drops                       | 5 ml = 1 tsp                           | 2 tabs               | 1 tab                     | ---              |
| 36-47 lbs    | 240 mg     |                                      | 7.5 ml = 1½ tsp                        | 3 tabs               | 1½ tab                    | ---              |
| 48-59 lbs    | 320 mg     |                                      | 10 ml = 2 tsp                          | 4 tabs               | 2 tabs                    | 1 tab            |
| 60-71 lbs    | 400 mg     |                                      | 12.5 ml = 2½ tsp                       | 5 tabs               | 2½ tabs                   | 1 tab            |
| 72-95 lbs    | 480 mg     |                                      | 15 ml = 3 tsp                          | 6 tabs               | 3 tabs                    | 1½ tab           |
| 96+ lbs      | 640 mg     |                                      | 20 ml = 4 tsp                          | 8 tabs               | 4 tabs                    | 2 tabs           |

## Benadryl (Diphenhydramine) Dosage Chart

**\*\* Benadryl can be given every 6 HOURS \*\* Consult your physician for children under 12 MOS OF AGE \*\***

| Weight (lbs) | Age (years) | Suspension (12.5mg/5ml) | Chewable (12.5 mg tab) | Fastmelts (12.5 mg tab) | Tab/Cap 25 mg |
|--------------|-------------|-------------------------|------------------------|-------------------------|---------------|
| 12-19 lbs    | 1-1½ yrs    | 2.5 ml = ½ tsp          |                        |                         |               |
| 20-26 lbs    | 1½-2 yrs    | 3.75 ml = ¾ tsp         |                        |                         |               |
| 27-39 lbs    | 2-4 yrs     | 5 ml = 1 tsp            | 1 tab                  | 1 tab                   |               |
| 40-52 lbs    | 5-6 yrs     | 7.5 ml = 1 ½ tsp        | 1 ½ tab                | 1 ½ tab                 |               |
| 53-67 lbs    | 7-8 yrs     | 10 ml = 2 tsp           | 2 tabs                 | 2 tabs                  | 1 tab/cap     |
| 68-79 lbs    | 9-10 yrs    | 10-12.5ml = 2-2½tsp     | 2-2½ tabs              | 2 tabs                  | 1 tab/cap     |
| 80-95 lbs    | 11-12 yrs   | 10-15 ml = 2-3 tsp      | 2-3 tabs               | 2-3 tabs                | 1 tab/cap     |
| 96+ lbs      | 12+ yrs     | 10-20 ml = 2-4 tsp      | 2-4 tabs               | 2-4 tabs                | 1-2 tabs/caps |

**\*\* CAUTION!!! Benadryl can cause significant sleepiness or an unexpected hyperactivity reaction in some children \*\***

**\*\* Dosing by weight is most accurate \*\* Consult your physician for any questions \*\***